

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____


IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.


This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims:

1. (original) A compound of formulae (I) or (II) having the structure



(I)



(II)

wherein

- $R_1, R_2, R_3, R_4, R_7, R_8, R_9, R_{10}, R_{11}, R_{12}, R_{14},$ and R_{15} are each, independently, hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16^-} , heteroaryl- R_{16^-} , hydroxyalkyl, HO- R_{16^-} , $R_{17-X-R_{16^-}}$, HS- R_{16^-} , $R_{17-S(O)-}$, $R_{17-S(O)_2-}$, R_{17-SO_3-} , $R_{17-S(O)_2NR-}$, $-N(R)_2$, $-NR-C(NH_2)=NR$, cyano, nitro, halogen, -OR, -SR, $-SO_3R$, $-S(O)_2N(R)_2$, $-C(O)R$, $-C(R)=N-OR$, $-C(NH_2)=NR$, $-CO_2R$, $-OC(O)R$, or $-C(O)N(R)_2$; or are taken together with either R_{p+1} or R_{p-1} linked with an -alkylene-, or -X-alkylene- group;
- R_5 is hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16^-} , heteroaryl- R_{16^-} , hydroxyalkyl, HO- R_{16^-} , $R_{17-X-R_{16^-}}$, HS- R_{16^-} , $-CR(O)$, $-CO_2R$, or $-C(O)N(R)_2$; or R_5 may be taken together with either R_6 or R_7 and linked with an -alkylene- or -X-alkylene- group;
- R_6 is hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16^-} , heteroaryl- R_{16^-} , hydroxyalkyl, HO- R_{16^-} , $R_{17-X-R_{16^-}}$, HS- R_{16^-} , $-CR(O)$, $-CO_2R$, or $-C(O)N(R)_2$; or R_6

may be taken together with either R₅ or R₇ and linked with an -alkylene- or -X-alkylene- group;

R₁₃ is R, R₁₇-X-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronidate;

R₁₆ is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;

R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;

R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;

X is O, -NR-, -S(O)_m-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

m is 0, 1, or 2;

p is 2, 3, 6, 7, 8, 9, 12, 13, or 14;

R₂₁, R₂₂, R₂₃, R₂₄, R₂₇, R₂₈, R₂₉, R₃₀, R₃₁, R₃₃, R₃₄, and R₃₅ are, independently, hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, R₁₇-SO₃-, R₁₇-S(O)₂NR-, -N(R)₂, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{q+1} or R_{q-1} linked with an -alkylene-, or -Y-alkylene- group;

R₂₅ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₅ may be taken together with either R₂₆ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₂₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₆ may be taken together with either R₂₅ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₃₂ is R, R₁₇-Y-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronidate;

Y is O, -NR-, -S(O)_n-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

n is 0, 1, or 2;

q is 22, 23, 26, 27, 28, 29, 32, 33, or 34;

or a pharmaceutically acceptable salt thereof.

2. (original) The compound according to claim 1, wherein the compound is of formula (I) or a pharmaceutical acceptable salt thereof.

3. (original) The compound according to claim 2, wherein R₁₃ is hydrogen, or a pharmaceutically acceptable salt thereof.

4. (original) The compound according to claim 3, wherein
R₁, R₂, R₃, R₄, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₄, and R₁₅ are each, independently, hydrogen, R₁₇,
aryl-R₁₆-, R₁₇-X-R₁₆-, hydroxyalkyl, HO-R₁₆-, halogen, -OR, -COR, or -CO₂R;
R₅ and R₆ are each, independently, hydrogen or R₁₇;
R₁₆ is -alkylene-;
R₁₇ is alkyl, aryl, heteroaryl, or perfluoroalkyl;
R is hydrogen or alkyl; or a pharmaceutically acceptable salt thereof.

5. (original) The compound according to claim 1, wherein the compound is of formula (II) or a pharmaceutical acceptable salt thereof.

6. (original) The compound according to claim 5, wherein R₃₂ is hydrogen, or a pharmaceutically acceptable salt thereof.

7. (original) The compound according to claim 6, wherein
R₂₁, R₂₂, R₂₃, R₂₄, R₂₇, R₂₈, R₂₉, R₃₀, R₃₁, R₃₃, R₃₄, and R₃₅ are each, independently, hydrogen,
R₁₇, aryl-R₁₆-, R₁₇-Y-R₁₆-, hydroxyalkyl, HO-R₁₆-, halogen, -OR, -COR, or -CO₂R;

R₂₅ and R₂₆ are each, independently, hydrogen or R₁₇;

R₁₆ is -alkylene-;

R₁₇ is alkyl, aryl, heteroaryl, or perfluoroalkyl;

R is hydrogen or alkyl; or a pharmaceutically acceptable salt thereof.

8. (currently amended) The compound according to claim 1, which is

- a) 4-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- b) 4-[(*S*)-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- c) 4-[(*R*)-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- d) 4-[(2-bromo-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- e) 2-methyl-4-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- f) 4-[(2-bromo-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]-2-methylphenol;
- g) 4-[(6-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- h) 4-[(2-bromo-6-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- i) 4-[(6-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- j) 4-[(*S*)-6-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- k) 4-[(*R*)-6-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- l) 4-[(2-bromo-6-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- m) 2-bromo-4-[(2-bromo-6-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- n) 4-[(6-*tert*-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- o) 4-[(*R*)-6-*tert*-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- p) 4-[(*S*)-6-*tert*-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- q) 4-[(2-bromo-6-*tert*-butylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- r) 4-[(6-ethylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- s) 4-[(2-bromo-6-ethylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- t) 4-[(6-ethylphenanthridin-5(6*H*)-yl)sulfonyl]-2-methylphenol;
- u) 4-[(2-bromo-6-ethylphenanthridin-5(6*H*)-yl)sulfonyl]-2-methylphenol;
- v) 4-[(*S*^{*})-6-[(*R*^{*})-1-methylpropyl]phenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- w) 4-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]benzene-1,2-diol;

- *) 2-hydroxy-5-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]benzoic acid;
- y) ethyl 2-ethoxy-5-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]benzoate;
- z) 2-(hydroxymethyl)-4-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- aa) 2-hydroxy-5-[(6-methylphenanthridin-5(6*H*)-yl)sulfonyl]benzaldehyde;
- bb) 4-[(6-ethyl-2-thien-3-ylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- cc) 4-[(6-ethyl-2-(3-methoxyphenyl)phenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- dd) 3-{6-ethyl-5-[(4-hydroxyphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}phenol;
- ee) 4-[(2-dibenzo[*b,d*]furan-4-yl-6-ethylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- ff) 4-[(8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- gg) 4-[(*S*)-8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- hh) 4-[(*R*)-8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- ii) 4-[(8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]-2-methylphenol;
- jj) 5-[(4-hydroxyphenyl)sulfonyl]-6-methyl-5,6-dihydrophenanthridin-9-ol;
- kk) 5-[(4-hydroxy-3-methylphenyl)sulfonyl]-6-methyl-5,6-dihydrophenanthridin-9-ol;
- ll) 5-[(4-hydroxy-3-methylphenyl)sulfonyl]-6-methyl-5,6-dihydrophenanthridin-7-ol;
- mm) 5-[(4-hydroxyphenyl)sulfonyl]-6-methyl-5,6-dihydrophenanthridin-7-ol;
- nn) 4-[(6-ethyl-8-fluorophenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- oo) 4-[(6-ethyl-8-fluorophenanthridin-5(6*H*)-yl)sulfonyl]-2-methylphenol;
- pp) 4-[(6-ethyl-7-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- qq) 4-[(6-ethyl-9-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- rr) 4-[(2-bromo-6-ethyl-8-fluorophenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- ss) 4-[(2-bromo-8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- tt) 2-chloro-4-[(6-ethyl-8-fluorophenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- uu) 4-[(6-ethyl-8-fluoro-2-phenylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- vv) 3-[(8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- ww) 2-fluoro-4-[(8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- xx) 4-[(8-fluoro-6-methylphenanthridin-5(6*H*)-yl)sulfonyl]benzene-1,2-diol;
- yy) 4-[(6-ethyl-8-fluoro-2-methylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- zz) 4-[(6-ethyl-8-fluoro-2-thien-3-ylphenanthridin-5(6*H*)-yl)sulfonyl]phenol;
- aaa) 4-[(6-ethyl-8-fluorophenanthridin-5(6*H*)-yl)sulfonyl]phenyl 3,3-dimethylbutanoate;


- bbb) 4-[(6-ethyl-8-fluorophenanthridin-5(6H)-yl)sulfonyl]phenyl propionate;
ccc) 4-[(6-ethyl-8-fluorophenanthridin-5(6H)-yl)sulfonyl]phenyl benzoate;
ddd) 2-fluoro-4-[(6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
eee) 4-[(2-bromo-6-methylphenanthridin-5(6H)-yl)sulfonyl]-2-fluorophenol;
fff) 4-[(6-methylphenanthridin-5(6H)-yl)sulfonyl]-2-(trifluoromethyl)phenol;
ggg) 2,6-dimethyl-4-[(6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
hhh) 4-[(6,8-dimethylphenanthridin-5(6H)-yl)sulfonyl]phenol;
iii) 4-[(8-chloro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
jjj) 4-[(2-bromo-8-chloro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
kkk) 2-{6-ethyl-5-[(4-hydroxyphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}phenol;
lll) 4-[[6-ethyl-2-[4-(methylthio)phenyl]phenanthridin-5(6H)-yl]sulfonyl]phenol;
mmm) 4-[[6-ethyl-2-[(E)-2-phenylethenyl]phenanthridin-5(6H)-yl]sulfonyl]phenol;
nnn) 4-[[2-(1,1'-biphenyl-4-yl)-6-ethylphenanthridin-5(6H)-yl]sulfonyl]phenol;
ooo) 4-[[2-(3-chlorophenyl)-6-ethylphenanthridin-5(6H)-yl]sulfonyl]phenol;
ppp) 4-[(6-ethyl-2-quinolin-8-ylphenanthridin-5(6H)-yl)sulfonyl]phenol;
qqq) 4-[(6-ethyl-2-phenylphenanthridin-5(6H)-yl)sulfonyl]phenol;
rrr) 4-[[6-ethyl-2-(2-methylphenyl)phenanthridin-5(6H)-yl]sulfonyl]phenol;
sss) 4-[(6-ethyl-2-thianthren-1-ylphenanthridin-5(6H)-yl)sulfonyl]phenol;
ttt) 4-[[2-(1-benzofuran-2-yl)-6-ethylphenanthridin-5(6H)-yl]sulfonyl]phenol;
uuu) 4-[[6-ethyl-2-(4-hydroxyphenyl)phenanthridin-5(6H)-yl]sulfonyl]phenol;
vvv) 4-[[2-(2-chlorophenyl)-6-ethylphenanthridin-5(6H)-yl]sulfonyl]phenol;
www) 4-[[6-ethyl-2-(4-ethylphenyl)phenanthridin-5(6H)-yl]sulfonyl]phenol;
xxx) 1-(5-{6-ethyl-5-[(4-hydroxyphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}thien-2-yl)ethanone;
yyy) 5-{6-ethyl-5-[(4-hydroxyphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}pyrimidine-2,4-diol;
zzz) 4-[[6-ethyl-2-(2-hydroxyphenyl)phenanthridin-5(6H)-yl]sulfonyl]-2-methylphenol;
aaaa) 4-[(6-ethyl-2-thien-3-ylphenanthridin-5(6H)-yl)sulfonyl]-2-methylphenol;
bbbb) 4-[[6-ethyl-2-[4-(methylthio)phenyl]phenanthridin-5(6H)-yl]sulfonyl]-2-methylphenol;

- eeee) 4-{{6-ethyl-2-[(E)-2-phenylethenyl]phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- dddd) 4-{{6-ethyl-5-[(4-hydroxy-3-methylphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}benzene-1,2-diol;
- eeee) 4-{{2-(1,1'-biphenyl-4-yl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- ffff) 4-{{6-ethyl-2-(3-hydroxyphenyl)phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- gggg) 4-{{2-(3-chlorophenyl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- hhhh) 4-{{6-ethyl-2-[(E)-hept-1-enyl]phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- iiii) 4-{{6-ethyl-2-pyridin-4-ylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- jjjj) 4-{{6-ethyl-2-quinolin-8-ylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- kkkk) 4-{{6-ethyl-2-(2-methylphenyl)phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- llll) 4-{{2-(1-benzothien-2-yl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- mmmm) 4-{{2-(1-benzothien-3-yl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- nnnn) 4-{{2-dibenzo[b,d]furan-4-yl-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- oooo) 4-{{2-(1-benzofuran-2-yl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- pppp) 4-{{6-ethyl-2-(4-hydroxyphenyl)phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- qqqq) 4-{{2-(2-chlorophenyl)-6-ethylphenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- rrrr) 4-{{6-ethyl-2-(4-ethylphenyl)phenanthridin-5(6H)-yl}sulfonyl}-2-methylphenol;
- ssss) 1-(5-{6-ethyl-5-[(4-hydroxy-3-methylphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}thien-2-yl)ethanone;
- tttt) 5-{6-ethyl-5-[(4-hydroxy-3-methylphenyl)sulfonyl]-5,6-dihydrophenanthridin-2-yl}pyrimidine-2,4-diol;
4-{{{(6R)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl}sulfonyl}phenol};
4-{{{(6S)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl}sulfonyl}phenol};

3-[[[(6R)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
3-[[[(6S)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
4-[[[(6S)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[[[(6R)-3,8-difluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[(3-fluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[(3-fluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[[[(6R)-3-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
3-[[[(6S)-3-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
4-[3-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonylbenzene-1,3-diol;
4-[[[(6R)-3-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[[[(6S)-3-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[(2-fluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[(2-fluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
4-[(3,9-difluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[(3,9-difluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
4-[(2,9-difluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[(2,9-difluoro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[(3-chloro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
3-[[[(6R)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
3-[[[(6S)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
4-[(3-chloro-6-methylphenanthridin-5(6H)-yl)sulfonyl]phenol;
4-[[[(6R)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
4-[[[(6S)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenol;
4-[[[(6R)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[[[(6S)-3-chloro-6-methylphenanthridin-5(6H)-yl]sulfonyl}benzene-1,3-diol;
4-[[[(6S)-8-fluoro-6-methylphenanthridin-5(6H)-yl]sulfonyl}phenyl sulfamate;
4-[(6-ethyl-8-fluoro-2-pyridin-3-ylphenanthridin-5(6H)-yl)sulfonyl]phenol;

or a pharmaceutically acceptable salt thereof.

9. (original) A pharmaceutical composition, which comprises a compound of formulae (I) or (II) having the structure



wherein

R₁, R₂, R₃, R₄, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₄, and R₁₅ are each, independently, hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, R₁₇-SO₃-, R₁₇-S(O)₂NR-, -N(R)₂, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{p+1} or R_{p-1} linked with an -alkylene-, or -X-alkylene- group;

R₅ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₅ may be taken together with either R₆ or R₇ and linked with an -alkylene- or -X-alkylene- group;

R₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₆ may be taken together with either R₅ or R₇ and linked with an -alkylene- or -X-alkylene- group;

R₁₃ is R, R₁₇-X-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronide;

R₁₆ is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;

R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;

R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;

X is O, -NR-, -S(O)_m-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

m is 0, 1, or 2;

p is 2, 3, 6, 7, 8, 9, 12, 13, or 14;

R₂₁, R₂₂, R₂₃, R₂₄, R₂₇, R₂₈, R₂₉, R₃₀, R₃₁, R₃₃, R₃₄, and R₃₅ are, independently, hydrogen, R₁₇; monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, R₁₇-SO₃-, R₁₇-S(O)₂N(R)₂-, -N(R)₂-, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂-, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{q+1} or R_{q-1} linked with an -alkylene-, or -Y-alkylene- group;

R₂₅ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₅ may be taken together with either R₂₆ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₂₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₆ may be taken together with either R₂₅ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₃₂ is R, R₁₇-Y-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronidate;

Y is O, -NR-, -S(O)_n-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

n is 0, 1, or 2;

q is 22, 23, 26, 27, 28, 29, 32, 33, or 34;

or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

10. (currently amended) A method of treating ~~or inhibiting~~ chronic inflammatory disease in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
11. (currently amended) A method of treating ~~or inhibiting~~ rheumatoid arthritis, spondyloarthropathies, osteoarthritis, psoriatic arthritis, or juvenile arthritis in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
12. (currently amended) A method of treating ~~or inhibiting~~ inflammatory bowel disease, Crohn's disease, ulcerative colitis, or indeterminate colitis in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
13. (currently amended) A method of treating ~~or inhibiting~~ psoriasis in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
14. (currently amended) A method of treating ~~or inhibiting~~ asthma or chronic obstructive pulmonary disease in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
15. (currently amended) A method of treating ~~or inhibiting~~ stroke, ischemia, or reperfusion injury in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.
16. (currently amended) A method of lowering cholesterol, triglycerides, Lp(a), and LDL levels; ~~inhibiting or~~ treating hypercholesteremia, hyperlipidemia, cardiovascular disease, atherosclerosis, acute coronary syndrome, peripheral vascular disease, restenosis, or vasospasm in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.

17. (currently amended) A method of treating ~~or inhibiting~~ Alzheimer's disease, cognitive decline, or senile dementia in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.


18. (currently amended) A method of treating ~~or inhibiting~~ type II diabetes in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.

19. (currently amended) A method of treating ~~or inhibiting~~ sepsis in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of claim 1.

20. (new) The compound according to claim 2, wherein R_{13} is $-S(O)_2NH_2$, or a pharmaceutically acceptable salt thereof.

21. (new) The compound according to claim 5, wherein R_{32} is $-S(O)_2NH_2$, or a pharmaceutically acceptable salt thereof.

22. (new) A process comprising providing a sulfonamide of formula 37:




wherein


R_1 , R_2 , R_3 , R_4 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{14} , and R_{15} are each, independently, hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} , heteroaryl- R_{16} , hydroxyalkyl,

- HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, R₁₇-SO₃-, R₁₇-S(O)₂NR-, -N(R)₂-, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂-, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{p+1} or R_{p-1} linked with an -alkylene-, or -X-alkylene- group;
- R₅ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₅ may be taken together with either R₆ or R₇ and linked with an -alkylene- or -X-alkylene- group;
- R₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-X-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₆ may be taken together with either R₅ or R₇ and linked with an -alkylene- or -X-alkylene- group;
- R₁₃ is R, R₁₇-X-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂-, or D-glucuronidate;
- R₁₆ is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;
- R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;
- R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;
- X is O, -NR-, -S(O)_m-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;
- m is 0, 1, or 2; and
- p is 2, 3, 6, 7, 8, 9, 12, 13, or 14; and

treating the sulfonamide of formula 37 with potassium carbonate to produce a phenanthridine of formula I:




23. (new) The process of claim 22 further comprising providing the S enantiomer of the biphenylamine of formula 36:



wherein


L is fluorine or chlorine; and

reacting the S enantiomer of the biphenylamine of formula 36 with a compound of formula 3 or an anhydride:




to produce a sulfonamide of formula 37.

24. (new) The process of claim 23 further comprising providing a biphenylamine of formula 36:



separating the biphenylamine of formula 36 into its respective enantiomers.


25. (new) The process of claim 24 further comprising providing a compound of formula 35:



reacting the compound of formula 35 with an ammonium source optionally in the presence of an acid catalyst to produce an intermediate imine; and

reducing the intermediate imine with a hydride source to produce a biphenylamine of formula 36.


26. (new) The process of claim 25 further comprising providing a compound of formula 33:



(33)

wherein

R₃₆ and R₃₇ are, independently, hydrogen or (C₁-C₄) lower straight chain or (C₃-C₆) branched chain alkyl, or R₃₆ and R₃₇ are taken together to form a pinacol moiety; and
reacting the compound of formula 33 in the presence of a coupling catalyst with a compound of formula 34:




(34)

wherein

W is a chlorine, bromine, or iodine atom, or a triflate (-OSO₂CF₃) moiety;
to produce a compound of formula 35.

27. (new) A process for preparing a compound of formula I:



(I)

wherein

$R_1, R_2, R_3, R_4, R_7, R_8, R_9, R_{10}, R_{11}, R_{12}, R_{14},$ and R_{15} are each, independently, hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} -, heteroaryl- R_{16} -, hydroxyalkyl, $HO-R_{16}$ -, $R_{17}-X-R_{16}$ -, $HS-R_{16}$ -, $R_{17}-S(O)$ -, $R_{17}-S(O)_2$ -, $R_{17}-SO_3$ -, $R_{17}-S(O)_2NR$ -, $-N(R)_2$ -, $-NR-C(NH_2)=NR$, cyano, nitro, halogen, $-OR$ -, $-SR$ -, $-SO_3R$ -, $-S(O)_2N(R)_2$ -, $-C(O)R$ -, $-C(R)=N-OR$ -, $-C(NH_2)=NR$ -, $-CO_2R$ -, $-OC(O)R$ -, or $-C(O)N(R)_2$; or are taken together with either R_{p+1} or R_{p-1} linked with an -alkylene-, or -X-alkylene- group;

R_5 is hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} -, heteroaryl- R_{16} -, hydroxyalkyl, $HO-R_{16}$ -, $R_{17}-X-R_{16}$ -, $HS-R_{16}$ -, $-CR(O)$ -, $-CO_2R$ -, or $-C(O)N(R)_2$; or R_5 may be taken together with either R_6 or R_7 and linked with an -alkylene- or -X-alkylene- group;

R_6 is hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} -, heteroaryl- R_{16} -, hydroxyalkyl, $HO-R_{16}$ -, $R_{17}-X-R_{16}$ -, $HS-R_{16}$ -, $-CR(O)$ -, $-CO_2R$ -, or $-C(O)N(R)_2$; or R_6 may be taken together with either R_5 or R_7 and linked with an -alkylene- or -X-alkylene- group;

R_{13} is R , $R_{17}-X-R_{16}$ -, $R_{17}-S(O)$ -, $R_{17}-S(O)_2$ -, $-SO_3R$ -, $-S(O)_2N(R)_2$ -, or D-glucuronidate;

R_{16} is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;

R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;

R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;


X is O, -NR-, -S(O)_m-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

m is 0, 1, or 2; and

p is 2, 3, 6, 7, 8, 9, 12, 13, or 14;

comprising

- reacting a compound of formula 33:




(33)

wherein

L is fluorine or chlorine; and

R₃₆ and R₃₇ are, independently, hydrogen or (C₁-C₄) lower straight chain or (C₃-C₆) branched chain alkyl, or R₃₆ and R₃₇ are taken together to form a pinacol moiety;

in the presence of a coupling catalyst with a compound of formula 34:




(34)

wherein


W is a chlorine, bromine, or iodine atom, or a triflate ($-\text{OSO}_2\text{CF}_3$) moiety;

to produce a compound of formula 35:




(35)

- reacting the compound of formula 35 with an ammonium source optionally in the presence of an acid catalyst to produce an intermediate imine;
- reducing the intermediate imine with a hydride source to produce a biphenylamine of formula 36:




(36)


- separating the biphenylamine of formula 36 into its respective enantiomers;
- reacting the S enantiomer of the biphenylamine of formula 36 with a compound of formula 3 or an anhydride:



to produce a sulfonamide of formula 37:




- treating the sulfonamide of formula 37 with potassium carbonate to produce a phenanthridine of formula I:



(II)

28. (new) A process comprising providing a sulfonamide of formula 37a:



(37a)

whercin

- R_{21} , R_{22} , R_{23} , R_{24} , R_{27} , R_{28} , R_{29} , R_{30} , R_{31} , R_{33} , R_{34} , and R_{35} are, independently, hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} -, heteroaryl- R_{16} -, hydroxyalkyl, HO- R_{16} -, R_{17} -Y- R_{16} -, HS- R_{16} -, R_{17} -S(O)-, R_{17} -S(O)₂-, R_{17} -SO₃-, R_{17} -S(O)₂NR-, -N(R)₂, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{q+1} or R_{q-1} linked with an -alkylene-, or -Y-alkylene- group;
- R_{25} is hydrogen, R_{17} , monofluoroalkyl, monofluoroalkenyl, aryl- R_{16} -, heteroaryl- R_{16} -, hydroxyalkyl, HO- R_{16} -, R_{17} -Y- R_{16} -, HS- R_{16} -, -CR(O), -CO₂R, or -C(O)N(R)₂; or R_{25}

may be taken together with either R₂₆ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₂₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O)-, -CO₂R, or -C(O)N(R)₂; or R₂₆ may be taken together with either R₂₅ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₃₂ is R, R₁₇-Y-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronidate;

R₁₆ is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;

R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;


R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;

Y is O, -NR-, -S(O)_n-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

n is 0, 1, or 2; and


q is 22, 23, 26, 27, 28, 29, 32, 33, or 34; and

treating the sulfonamide of formula 37a with potassium carbonate to produce a phenanthridine of formula II:



(II)

29. (new) The process of claim 28 further comprising providing the S enantiomer of the biphenylamine of formula 36a:




(36a)

wherein

L is fluorine or chlorine; and


reacting the S enantiomer of the biphenylamine of formula 36a with a compound of formula 32 or an anhydride:



(32)


to produce a sulfonamide of formula 37a.

30. (new) The process of claim 29 further comprising providing a biphenylamine of formula 36a:



separating the biphenylamine of formula 36a into its respective enantiomers.


31. (new) The process of claim 30 further comprising providing a compound of formula 35a:



reacting the compound of formula 35a with an ammonium source optionally in the presence of an acid catalyst to produce an intermediate imine; and


reducing the intermediate imine with a hydride source to produce a biphenylamine of formula 36.

32. (new) The process of claim 31 further comprising providing a compound of formula 33a:



wherein


R₃₆ and R₃₇ are, independently, hydrogen or (C₁-C₄) lower straight chain or (C₃-C₆) branched chain alkyl, or R₃₆ and R₃₇ are taken together to form a pinacol moiety; and
reacting the compound of formula 33a in the presence of a coupling catalyst with a compound of formula 34a:



wherein

W is a chlorine, bromine, or iodine atom, or a triflate (-OSO₂CF₃) moiety;
to produce a compound of formula 35.

33. (new) A process for preparing a compound of formula II:



wherein

R₂₁, R₂₂, R₂₃, R₂₄, R₂₇, R₂₈, R₂₉, R₃₀, R₃₁, R₃₃, R₃₄, and R₃₅ are, independently, hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, R₁₇-SO₃-, R₁₇-S(O)₂NR-, -N(R)₂, -NR-C(NH₂)=NR, cyano, nitro, halogen, -OR, -SR, -SO₃R, -S(O)₂N(R)₂, -C(O)R, -C(R)=N-OR, -C(NH₂)=NR, -CO₂R, -OC(O)R, or -C(O)N(R)₂; or are taken together with either R_{q+1} or R_{q-1} linked with an -alkylene-, or -Y-alkylene- group;

R₂₅ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₅ may be taken together with either R₂₆ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₂₆ is hydrogen, R₁₇, monofluoroalkyl, monofluoroalkenyl, aryl-R₁₆-, heteroaryl-R₁₆-, hydroxyalkyl, HO-R₁₆-, R₁₇-Y-R₁₆-, HS-R₁₆-, -CR(O), -CO₂R, or -C(O)N(R)₂; or R₂₆ may be taken together with either R₂₅ or R₂₇ and linked with an -alkylene- or -Y-alkylene- group;

R₃₂ is R, R₁₇-Y-R₁₆-, R₁₇-S(O)-, R₁₇-S(O)₂-, -SO₃R, -S(O)₂N(R)₂, or D-glucuronidate;

R₁₆ is -alkylene-, -cycloalkylene-, -alkylene-X-alkylene-, -alkylene-X-cycloalkylene-, -cycloalkylene-X-alkylene-, or -cycloalkylene-X-cycloalkylene-;

R₁₇ is alkyl, aryl, heteroaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkenyl-X-alkylene-, cycloalkenyl-X-alkylene-, or perfluoroalkyl;

R is, independently, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, monofluoroalkyl, perfluoroalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy-(C₂-C₆)alkyl, alkoxyalkyl, alkylthioalkyl, formyl, acyl, alkoxycarbonyl, -C(O)NH₂, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminoalkyl, or dialkylaminoalkyl; or when an atom contains two R groups, the R groups may be taken together linked with an -alkylene- group;


Y is O, -NR-, -S(O)_n-, -C(O)-, -OC(O)-, -C(O)O-, -NRC(O)-, or -C(O)NR-;

n is 0, 1, or 2;

q is 22, 23, 26, 27, 28, 29, 32, 33, or 34;

comprising

- reacting a compound of formula 33a:




(33a)

wherein

L is fluorine or chlorine; and

R₃₆ and R₃₇ are, independently, hydrogen or (C₁-C₄) lower straight chain or (C₃-C₆) branched chain alkyl, or R₃₆ and R₃₇ are taken together to form a pinacol moiety;

in the presence of a coupling catalyst with a compound of formula 34a:




(34a)


wherein

W is a chlorine, bromine, or iodine atom, or a triflate (-OSO₂CF₃) moiety;


to produce a compound of formula 35a:




- reacting the compound of formula 35a with an ammonium source optionally in the presence of an acid catalyst to produce an intermediate imine;
- reducing the intermediate imine with a hydride source to produce a biphenylamine of formula 36a:



- separating the biphenyl amine of formula 36a into its respective enantiomers;
- reacting the S enantiomer of the biphenylamine of formula 36a with a compound of formula 32 or an anhydride:



to produce a sulfonamide of formula 37a:



; and

- treating the sulfonamide of formula 37a with potassium carbonate to produce a phenanthridine of formula II:

